

# **Utah Weekly Pertussis Update**

2016 data, through MMWR week ending January 23, 2016 (MMWR week 3)


Data presented in this report are based on current data available and will be subject to change weekly.

### **Data Summary**

Total number of cases reported, 2016 year-to-date		Utah Incidence rate, 2016 year-to-date (YTD) per
(YTD)	15	100,000 person-years
		Utah Incidence rate, 2015 per 100,000 person-
Total number of cases reported 2015	483	years
		National Incidence rate, 2014 per 100,000 person-
Number of cases reported in past week	8	years.
Total number of cases reported, through same time		
period 2015	39	2016 Hospitalizations
	<1 year	
Age groups with the highest rates	1-4 year	2016 Cases in infants <1 year

<sup>\*</sup>National Incidence Rate is updated as released by the Centers for Disease Control and Prevention.

## Number of cases reported by week



<sup>\*</sup>Additional cases may have occurred, especially in the most recent 3 weeks that have not yet been reported, as indicated by the grayed out section of the graph above.

### 2016 Year-to-Date Incidence Rates, By Age

Age	2016	2014	
(years)	Cases	population	Rate*, **
<1	2	50,629	4.7
1-4	3	201,502	1.8
5-14	3	511,355	0.7
15-24	3	474,017	0.8
25-34	1	440,622	0.3
35-44	0	386,350	0.0
45-54	2	306,598	0.8
55-64	1	276,569	0.4
65+	0	295,260	0.0
Total	15	2,942,902	13.3

<sup>\*</sup> Rate is calculated per 100,000 person years.

Infants less than 1 year of age have historically had the highest incidence rates of pertussis due to their susceptibility.

School Age Incidence Rates, 2016 Year-to-Date


		Age		2014	
	Grade	(years)	<b>2016 Cases</b>	Population	Rate*
	Preschool	3-4	6	101,497	7.1
Kii	ndergarten	5	0	52,285	0.0
	1	6	1	53,101	2.3
	2	7	0	52,876	0.0
	3	8	1	52,082	2.3
	4	9	0	51,852	0.0
	5	10	0	51,206	0.0
	6	11	0	50,689	0.0
	7	12	0	48,996	0.0
	8	13	0	49,110	0.0
	9	14	1	49,158	2.4
	10	15	1	47,568	2.5
	11	16	0	46,830	0.0
	12	17	2	46,231	5.2

<sup>\*</sup> Rate is calculated per 100,000 person-years

School age incidence rates were calculated using the average age of a student in each grade.

The high incidence rates in 5th and 6th grade may be due to waining immunity.

**Utah Case Counts of Pertussis, 2011-2016\* Year to Date** 



Pertussis tends to be cyclical. Reports have increased in Utah since 2010, peaked in 2012, and are currently trending downward.


\* 2016 data includes pertussis cases reported year to date. 2010-2014 data reflects cases reported during the entire calendar year for each year.



## Incidence of pertussis by Local Health Department, 2016 YTD

	2016	
Local Health Department	Cases	Rate*
Bear River	1	0.7
Central Utah	0	0.0
Davis County	0	0.0
Salt Lake Valley	4	0.4
San Juan County	0	0.0
Southeast Utah	0	0.0
Southwest Utah	0	0.0
<b>Summit County</b>	0	0.0
<b>Tooele County</b>	0	0.0
TriCounty	0	0.0
Utah County	6	1.3
Wasatch County	0	0.0
Weber-Morgan	4	1.9

<sup>\*</sup> Rate is calculated per 100,000 person-years using 2014 Utah population estimates





13.3	
16.7	
10.4	
0	
2	



Э

er 100,000

r 100,000

r 100,000

r 100,000

100,000